

# MathQuest: Differential Equations

## Equilibria and Stability

1. The differential equation  $\frac{dy}{dt} = (t - 3)(y - 2)$  has equilibrium values of
- (a)  $y = 2$  only
  - (b)  $t = 3$  only
  - (c)  $y = 2$  and  $t = 3$
  - (d) No equilibrium values

*Answer: (a).* An equilibrium value is a value of the function  $y$ , that makes the derivative  $\frac{dy}{dt} = 0$  for all  $t$ , so if we choose this as our initial condition, then the function will always remain at this value.

CC KC MA232 S07: **0**/11/89/0 time 1:45

CC MP MA334 S07: **15**/8/77/0

CC HZ MA232 S08: **4**/0/96/0 time 1:10 revote: **56**/37/0/7 time 2:30

HHS JG MA232 S08: **0**/0/100/0

AU CS MA244 S08: **0**/0/100/0

CC JS MA232 S09: **44**/17/39/0

by Carroll College MathQuest

DEQ.00.05.005

2. Suppose that 3 is an equilibrium value of a differential equation. This means that
- (a) the values will approach 3.
  - (b) if the initial value is below 3, the values will decrease.
  - (c) if the initial value is 3, then all of the values will be 3.
  - (d) all of the above.

*Answer: (c).* If we choose an equilibrium value as our initial condition, this means that the derivative will be zero, so we will stay at this value.

CC KC MA232 S07: 0/0/**90**/10 time 1:25

CC MP MA334 S07: 12/4/**65**/19

WH GC MAT345 S08: 33/67/**0**/0

HHS JG MA232 S08: 0/0/**100**/0

AU CS MA244 S08: 41/4/**55**/0

CC JS MA232 S09: 0/0/**86**/14

by Carroll College MathQuest

DEQ.00.05.008

3. We know that a given differential equation is in the form  $y' = f(y)$ , where  $f$  is a continuous function of  $y$ . Suppose that  $f(5) = 2$  and  $f(-1) = -6$ .
- (a)  $y$  must have an equilibrium value between  $y = 5$  and  $y = -1$ .
  - (b)  $y$  must have an equilibrium value between  $y = 2$  and  $y = -6$ .
  - (c) This does not necessarily indicate that any equilibrium value exists.

*Answer: (a).* The function  $f(y)$  gives us the derivative of  $y$  and if it is continuous, this means that at some point between  $y = 5$  and  $y = -1$  there must be a value where  $y' = f(y) = 0$ , and that makes it an equilibrium value.

CC KC MA232 S07: **33**/10/57 time 1:45

WH GC MAT345 S08: **57**/43/0

CC HZ MA232 S08: **26**/15/59 time 2:50

HHS JG MA232 S08: **0**/0/100

by Carroll College MathQuest

DEQ.00.05.010

4. We know that a given differential equation is in the form  $y' = f(y)$ , where  $f$  is a continuous function of  $y$ . Suppose that  $f(10) = 0$ ,  $f(9) = 3$ , and  $f(11) = -3$ .
- (a) This means that  $y = 10$  is a stable equilibrium.
  - (b)  $y = 10$  is an equilibrium, but it might not be stable.
  - (c) This does not tell us for certain that  $y = 10$  is an equilibrium.

*Answer: (b).*  $f(9) = 3$  means that when  $y = 9$  the function is increasing, and  $f(11) = -3$  means that when  $y = 11$  the function is decreasing. This might lead us to believe that solutions must go towards the equilibrium at  $y = 10$ , making it a stable equilibrium. However since we don't know the function, we can't be sure this is true.  $y = 10$  might be unstable, with two stable equilibria located at  $y = 9.5$  and  $y = 10.5$ .

CC KC MA232 S07: 86/**14**/0 time 1:30

CC HZ MA232 S08: 67/**33**/0 time 1:45

HHS JG MA232 S08: 0/**58**/42

by Carroll College MathQuest

DEQ.00.05.020

5. We know that a given differential equation is in the form  $y' = f(y)$ , where  $f$  is a continuous function of  $y$ . Suppose that  $f(6) = 0$ ,  $f(14) = 0$ , and  $y(10) = 10$ .
- (a) This means that  $y(0)$  must have been between 6 and 14.
  - (b) This means that  $y(20) = 0$  is impossible.
  - (c) This means that  $y(20) = 20$  is impossible.
  - (d) All of the above.
  - (e) None of the above.

*Answer: (d).* A solution trajectory can never cross an equilibrium solution trajectory, so if  $y(10) = 10$ , this means that all past and future values of  $y$  are confined between the equilibria at  $y = 6$  and  $y = 14$ .

CC KC MA232 S07: 14/5/5/0/76 time 2:30

CC HZ MA232 S08: 0/0/4/0/96 time 2:20

HHS JG MA232 S08: 0/0/17/0/83

AU CS MA244 S08: 17/22/6/11/44

by Carroll College MathQuest

DEQ.00.05.030

6. We know that a given differential equation is in the form  $y' = f(y)$ , where  $f$  is a continuous function of  $y$ . Suppose that  $f(2) = 3$  and that  $y(0) = 0$ . Which of the following is impossible?
- (a)  $y(10) = 6$
  - (b)  $y(10) = -6$
  - (c)  $y(-10) = 6$
  - (d)  $y(-10) = -6$
  - (e) All of these are possible

*Answer: (c).* When  $y = 2$ , we know that  $y' = 3$  so the function  $y(t)$  is increasing. If  $y(-10) = 6$ , then for  $t > -10$ ,  $y$  must always be larger than  $y = 2$ : If it went down through  $y = 2$ , then  $y'$  would have to be negative.

WH GC MAT345 S08: 0/0/14/0/86

HHS JG MA232 S08: 0/8/0/0/92

by Carroll College MathQuest

DEQ.00.05.040

7. We know that a given differential equation is in the form  $y' = f(y)$ , where  $f$  is a continuous function of  $y$ . Suppose that  $f(5) = -2$ ,  $f(10) = 4$ , and that  $y(10) = 3$ .
- (a)  $y(0)$  must be below 5.
  - (b)  $y(20)$  must be below 5.
  - (c)  $y(5)$  could be above 10.
  - (d)  $y(15)$  must be less than 3.

*Answer: (b).*  $y(20)$  must be below 5, because we know that  $y$  is decreasing when  $y = 5$ . Because  $y(10) = 3$  this means that in the future,  $y$  could never rise above 5. (a) is not true because we know that the function is decreasing when  $y = 5$ . Thus  $y$  could have started out above 5, and then decreased to 3 when  $t = 10$ . (c) is not true because we know that the function is increasing when  $y = 10$ , so if  $y(5)$  was greater than 10, then  $y$  would always have to be greater than 10 for  $y > 5$ . (d) is not true because we don't know what  $f(3)$  is. If  $f(3)$  was positive, then  $y(15)$  could be larger than 3. We can only say for certain that it must be below  $y = 5$ .

HHS JG MA232 S08: 0/75/25/0

by Carroll College MathQuest

DEQ.00.05.050

8. A differential equation has a stable equilibrium value of  $T = 6$ . Which of the following functions is definitely not a solution?
- (a)  $T(t) = 5e^{-3t} + 6$
  - (b)  $T(t) = -4e^{-2t} + 6$
  - (c)  $T(t) = 4e^{2t} + 10$
  - (d) They could all be solutions

*Answer: (d).* Both a and b asymptotically approach 6. c diverges from 10, but just because 6 is a stable equilibrium doesn't mean that the equation does not also have an unstable equilibrium at 10.

CC KC MA232 S07: 22/6/66/16 time 4:30

HHS JG MA232 S08: 33/0/58/8

by Carroll College MathQuest

DEQ.00.05.060

9. Consider the differential equation  $\frac{df}{dx} = \sin(f)$
- (a)  $f = 0$  is a stable equilibrium.

- (b)  $f = 0$  is an unstable equilibrium.
- (c)  $f = 0$  is not an equilibrium.

*Answer:* (b).  $f = 0$  is an equilibrium, because it makes the derivative  $\frac{df}{dx} = 0$ . This equilibrium is unstable because initial conditions slightly above this equilibrium will give us positive derivatives, pushing the solution trajectory away from equilibrium, while conversely initial conditions slightly below equilibrium give us negative derivatives, also pushing solutions away from equilibrium.

CC KC MA232 S07: 24/48/29 time 2:00  
CC HZ MA232 S08: 38/62/0 time 2:10 Review  
AU CS MA244 S08: 10/74/16  
CC JS MA232 S09: 40/50/10

by Carroll College MathQuest  
DEQ.00.05.070


10. Consider the differential equation  $\frac{df}{dx} = af + b$ , where  $a$  and  $b$  are positive parameters. If we increase  $b$ , what will happen to the equilibrium value?
- (a) it increases
  - (b) it decreases
  - (c) it stays the same
  - (d) not enough information is given

*Answer:* (b). The equilibrium value of this equation is  $-b/a$ , so if  $b$  becomes larger, this negative quantity decreases.

CC MP MA334 S07: 19/35/27/19

by Carroll College MathQuest  
DEQ.00.05.080

11. Suppose that  $\frac{dy}{dt} = f(y)$ , which is plotted below. What are the equilibrium values of the system?



- (a)  $y = \frac{1}{2}$  is the only equilibrium.
- (b)  $y = -1$  and  $y = 2$  are both equilibria.
- (c) Not enough information is given.

*Answer: (b).* According to the graph  $y' = f(y) = 0$  when  $y = -1$  and  $y = 2$ , so these are equilibria.

CC KC MA232 S07: 5/76/19 time 1:40

WH GC MAT345 S08: 86/14/0

CC HZ MA232 S08: 15/63/22 time 2:50

HHS JG MA232 S08: 9/91/0


AU CS MA244 S08: 0/92/8

CC JS MA232 S09: 19/67/14

by Carroll College MathQuest

DEQ.00.05.083

12. Suppose that  $\frac{dy}{dt} = f(y)$ , which is plotted below. What can we say about the equilibria of this system?



- (a)  $y = 0$  is stable,  $y = \pm 2$  are unstable.
- (b)  $y = 0$  is unstable,  $y = \pm 2$  are stable.
- (c)  $y = -2, 0$  are stable,  $y = 2$  is unstable.
- (d)  $y = -2$  is unstable,  $y = 0, 2$  are unstable
- (e) None of the above

*Answer: (a).* Equilibria are points where  $y' = f(y) = 0$  so the equation has equilibria at  $y = 0, \pm 2$ . A stable equilibrium is a point where  $f$  changes sign from positive to negative, so  $y = 0$  is stable, while conversely  $y = \pm 2$  are unstable.

CC KC MA232 S07: **60**/0/5/15/20 time 2:00

WH GC MAT345 S08: **100**/0/0/0/0

CC HZ MA232 S08: **36**/45/9/0/9 time 3:15 Review

HHS JG MA232 S08: **36**/45/9/0/9

AU CS MA244 S08: **63**/13/0/0/25

CC JS MA232 S09 **86**/0/0/14/0

by Carroll College MathQuest

DEQ.00.05.085

13. **True or False** Could a differential equation have infinitely many equilibria?

*Answer: (True).* For example if  $\frac{dy}{dx} = \sin y$  then equilibria exists at  $y = n\pi$  for any integer  $n$ .

by Carroll College MathQuest

DEQ.00.05.090

14. **True or False** Could a differential equation have infinitely many equilibria over a finite range?

*Answer: (True).* For example if  $\frac{dy}{dx} = 0$  then all values of  $y$  are equilibria.

by Carroll College MathQuest

DEQ.00.05.100

15. Consider the differential equation  $\frac{df}{dx} = af + b$ , where  $a$  and  $b$  are non-negative parameters. This equation would have no equilibrium if

- (a)  $a = 0$
- (b)  $b = 0$
- (c)  $a = 1$
- (d) More than one of the above

*Answer: (a).* If  $a = 0$  the differential equation is just  $\frac{df}{dx} = b$  where  $b$  is a positive number. This will result in an increasing function, so there can be no equilibrium.

by Carroll College MathQuest

DEQ.00.05.110

16. What is the equilibrium value of  $\frac{dg}{dz} = -\frac{1}{2}g + 3e^z$ ?


- (a) This system is at equilibrium when  $g = 6e^z$ .
- (b) This system is at equilibrium when  $z = \ln\left(\frac{g}{6}\right)$ .
- (c) Both a and b are true.
- (d) This equation has no equilibrium.

*Answer: (d).* An equilibrium would be a value of the function  $g$ , that would make  $g' = 0$  for all values of  $z$ . In this case, there is no such value.

by Carroll College MathQuest

DEQ.00.05.120

17. The figure below plots several functions which all solve the differential equation  $y' = ay + b$ . What could be the values of  $a$  and  $b$ ?



- (a)  $a = 1, b = 3$
- (b)  $a = 2, b = -6$
- (c)  $a = -1, b = -3$
- (d)  $a = -2, b = 6$
- (e)  $b = 3$  but  $a$  is not easy to tell

*Answer: (d).* The plot shows the values leveling off at  $y = 3$ , so the system has a stable equilibrium at  $y = 3$ . Because the equilibrium is stable, this means that  $a < 0$ . To be an equilibrium value we need  $-\frac{b}{a} = 3$ .

CC KC MA232 S07: 0/24/24/48/5 time 3:30


CC HZ MA232 S08: 0/24/3/0/72 time 2:10 Review

HHG JG MA232 S08: 9/9/0/27/55

by Carroll College MathQuest

DEQ.00.05.130

18. The figure below plots several functions which all solve the differential equation  $\frac{dy}{dx} = ay + b$ . What could be the values of  $a$  and  $b$ ?



- (a)  $a = 0.5, b = 2$
- (b)  $a = 0.5, b = -2$
- (c)  $a = -0.5, b = 2$
- (d)  $a = -0.5, b = -2$
- (e) None of the above are possible.

*Answer:* (a). The plot shows an unstable equilibrium at  $y = -4$ . This means that  $a$  must be positive and  $y_{eq} = -\frac{b}{a} = -4$ , which is satisfied if  $a = 0.5$  and  $b = 2$ .

WH GC MAT345 S08: **67**/17/17/0/0

Carroll College MathQuest

DEQ.00.05.140